

---

---

## Integrated Stratigraphy of the Austin Chalk Group across the San Marcos Arch, Bexar County, Texas

John Cooper, Jeffrey Hardwick, and Alexis Godet

Department of Geological Sciences, University of Texas at San Antonio,  
1 UTSA Circle, San Antonio, Texas 78249

GCAGS Explore & Discover Article #00057\*

[http://www.gcags.org/exploreanddiscover/2016/00057\\_cooper\\_et\\_al.pdf](http://www.gcags.org/exploreanddiscover/2016/00057_cooper_et_al.pdf)

Posted September 13, 2016.

\*Article based on an extended abstract published in the *GCAGS Transactions* (see footnote reference below), which is available as part of the entire 2016 *GCAGS Transactions* volume via the GCAGS Bookstore at the Bureau of Economic Geology ([www.beg.utexas.edu](http://www.beg.utexas.edu)) or as an individual document via AAPG Datapages, Inc. ([www.datapages.com](http://www.datapages.com)), and delivered a poster presentation at the 66th Annual GCAGS Convention and 63rd Annual GCSSEPM Meeting in Corpus Christi, Texas, September 18–20, 2016.

---

---

### EXTENDED ABSTRACT

The Cretaceous Period was punctuated by a series of perturbations of the global carbon cycle that culminated in the installation of oceanic anoxia and affected marine ecosystems (Schlanger and Jenkyns, 1976; Weissert et al., 1998; Erba, 2004; Weissert and Erba, 2004). One of these oceanic anoxic events (OAE), the OAE3, developed during the Coniacian to early Campanian time period and lasted for almost 5 Myr, although its extent is regional compared to other OAEs, such as the OAE2 at the Cenomanian – Turonian boundary (Wagner et al., 2004; Locklair et al., 2011; Wagreich, 2012). The OAE3 is depicted by a series of 5 short-amplitude, high-frequency positive peaks in the evolution of carbon stable isotope ( $\delta^{13}\text{C}$ ; Wagreich, 2009). During this part of the Late Cretaceous, the epicontinental Western Interior Seaway covered central Texas, resulting in the deposition of a thick series of fine-grained carbonate sediments known as the Austin Chalk. Despite a relatively well-constrained biostratigraphic framework (e.g., Young, 1963; Pessagno, 1969; Jiang, 1989; Lundquist, 2000), the Austin Chalk lacks a robust integrated stratigraphic framework in which the impact of the aforementioned paleoenvironmental crisis as well as more local parameters (e.g., local submarine topography, nutrient and detrital input) on carbonate bio- and lithofacies could be assessed.

...



