
Sediment Dispersal Patterns of the Outer Shelf to Upper Slope Paleocene-Eocene Wilcox Group, South-Central Texas Coast

Hongliu Zeng¹, William A. Ambrose¹, and Wenlong Xu²

¹Bureau of Economic Geology, Jackson School of Geosciences, University of Texas at Austin, University Station, Box X, Austin, Texas 78713–8924, U.S.A.

²Excellong, Inc., 77 Sugar Creek Center Blvd., Ste. 215, Sugar Land, Texas 77478, U.S.A.

GCAGS Explore & Discover Article #00183*

http://www.gcags.org/exploreanddiscover/2016/00183_zeng_et_al.pdf

Posted September 13, 2016.

*Abstract extracted from a full paper published in the *GCAGS Journal* (see footnote reference below), which is available as part of the entire 2016 *GCAGS Journal* volume via the GCAGS Bookstore at the Bureau of Economic Geology (www.beg.utexas.edu) or as an individual open-access document via www.gcags.org.

ABSTRACT

Applying wavelet-phase adjustment and attribute analysis to interpret lithology and stratal slices to deduce seismic geomorphology, we interpreted a 960 mi² (2500 km²) 3D seismic volume, wireline log, and sparse-core data along the south-central Texas coast to investigate the sediment dispersal patterns in transition from on-shelf to the deepwater. The sequence-stratigraphic architecture of the Wilcox Group revealed four subbasins separated by major growth faults, which roughly correlated to a lower Wilcox subunit, a middle Wilcox subunit, and two upper Wilcox subunits. Each of the subbasins developed over an average 3.5 m.y. period. We prepared lithology-calibrated stratal slice maps to reveal high-resolution (33–130-ft [10–40 m]) sediment-dispersal patterns and associated systems tracts in subbasin C of the upper Wilcox. Recognized depositional systems include: (1) incised valley fills (IVFs) and relict on-shelf deltas on an exposed shelf; (2) off-shelf lowstand prograding deltaic systems composed of lobate deltaic sandstone bodies; and (3) off-shelf slope fans best characterized by point-source, fanlike channel/levee systems. At least four episodes of shallow LST (lowstand system tract) incised valleys containing 33–130 ft (10–40 m) of gross sandstone inferred from wireline logs were identified. These thin, sandstone-rich IVFs on exposed shelf are coeval with thick, sand–shale mixed slope fans and LST deltaic systems along growth faults, suggesting that large amounts of sediment may have bypassed these channels and may have been trapped between the fault zones in multiple subbasins. In addition to deep-cut submarine canyons (e.g., Yoakum Canyon), these IVFs provided an alternative fairway that may have transported sandy LST deposits and associated reservoirs seaward of the Cretaceous shelf edge to the deepwater Gulf of Mexico.

Originally published as: Zeng, H., W. A. Ambrose, and W. Xu, 2016, Sediment dispersal patterns of the outer shelf to upper slope Paleocene-Eocene Wilcox Group, south-central Texas Coast: *Gulf Coast Association of Geological Societies Journal*, v. 5, p. 215–237.