
Seismic Characterization of the Upper Cretaceous Submarine Fan System in the Deepwater Foz do Amazonas Basin, Northern Brazil: An analog for Upper Cretaceous Fan Plays on Atlantic Passive Margins

Lucia Torrado

Department of Earth and Atmospheric Sciences, University of Houston,
Science & Research Bldg. 1, Rm. 312, Houston, Texas 77204-5007

GCAGS Explore & Discover Article #00286*
http://www.gcags.org/exploreanddiscover/2017/00286_torrado.pdf
Posted October 30, 2017.

* Article based on an abstract published in the *GCAGS Transactions* (see footnote reference below), which is available as part of the entire 2017 *GCAGS Transactions* volume via the GCAGS Bookstore at the Bureau of Economic Geology (www.beg.utexas.edu) or as an individual document via AAPG Datapages, Inc. (www.datapages.com), and delivered as a poster presentation at the 67th Annual GCAGS Convention and 64th Annual GCSSEPM Meeting in San Antonio, Texas, November 1–3, 2017.

ABSTRACT

The deepwater Foz do Amazonas Basin shares the same working, petroleum system linked to the 2011 Zaedyus Field (700 million barrels of oil) and 2017 Liza Field (800 million barrels of oil equivalent) discovered in the French Guiana–Guyana–Suriname basins to the northwest: A Cenomanian-Turonian age source rock charging Upper Cretaceous deepwater fan reservoirs. We conducted seismic facies analysis and geomorphology on the Upper Cretaceous submarine fan systems (Limoeiro Formation) using 21,369 km of 2D depth-converted seismic tied to 3 exploration wells, in order to: (1) distinguish mud-prone vs. sand-prone deposits on seismic data; (2) understand depositional controls on stratigraphic traps within the fans; and (3) determine reservoir quality of fan sandstones. The Upper Cretaceous deepwater fan system of the Foz do Amazonas Basin deposited ~40 km seaward from the paleoshelf break and includes: (1) upper slope fans interpreted as sand-prone, channel-levee complexes expressed on reflection data as concave, discontinuous reflections with moderate amplitude strength; (2) basin floor fans seen as high-amplitude, continuous reflection packages (HARPs) of sand sheets with little to no structural deformation; (3) shingled turbidites expressed as offlapping reflections of moderate amplitude strength, and (4) shale packages of low-amplitude strength reflections. These large Upper Cretaceous fan complexes (~800 km²) were sourced by mature terrigenous sands eroded from hills located close to the paleo-shoreline, and deposited in conditions of: (1) ~3.19° gradient along a ~95 km long slope; (2) strong turbidity currents; and (3) steepening of the margin as a result of rifting and passive margin subsidence. These factors combine to create conditions for good reservoir quality and trap preservation with multiple seals and traps.

Originally published as: Torrado, L., 2017, Seismic characterization of the Upper Cretaceous submarine fan system in the deepwater Foz do Amazonas Basin, northern Brazil: An analog for Upper Cretaceous fan plays on Atlantic passive margins: Gulf Coast Association of Geological Societies Transactions, v. 67, p. 651.