
High-Resolution Facies Analysis and Depositional Architecture of the Buda Formation in Central Texas

Xiaodong Zhang and Michael C. Pope

Department of Geology and Geophysics, Texas A&M University,
3115 TAMU, College Station, Texas 77843-3115

GCAGS Explore & Discover Article #00293*

http://www.gcags.org/exploreanddiscover/2017/00293_zhang_and_pope.pdf

Posted October 30, 2017.

*Article based on an abstract published in the *GCAGS Transactions* (see footnote reference below), which is available as part of the entire 2017 *GCAGS Transactions* volume via the GCAGS Bookstore at the Bureau of Economic Geology (www.beg.utexas.edu) or as an individual document via AAPG Datapages, Inc. (www.datapages.com), and delivered as a poster presentation at the 67th Annual GCAGS Convention and 64th Annual GCSSEPM Meeting in San Antonio, Texas, November 1-3, 2017.

ABSTRACT

The Upper Cretaceous Buda Formation outcrop extends from West Texas to northern Fort Worth County, and extending continuously in the subsurface from South to Central Texas. Three facies (from deepest to shallowest) are interpreted from core and outcrops: foraminifera mudstone, skeletal wackestone/packstone, and massive mudstone. Outcrops are mainly composed of skeletal wackestone with massive mudstone, while cores are dominated by globigerinid mudstone and skeletal wackestone/packstone. Both outcrops and cores are extremely burrow mottled. 2D strike and dip cross sections constrained by cores indicate the thickness of the Buda Formation varies significantly from 20 to 100 feet through Central Texas. Considerable thickening of the Buda Formation within the trough between Edwards and Sligo paleo-shelf margins occurs in southern Wilson, northern Karnes, and Gonzales counties of Central Texas. The dramatic variations of the Buda Formation thickness along the dip direction are consistent with filling differential accommodation space across the paleotopographic profile of the pre-existing Lower Cretaceous reef trend. Meanwhile, facies variations from the outcrop and subsurface data record a transitional change of the depositional environments from inner ramp to subtidal shallow marine. The abundance of bioturbation and benthic forams across the depositional profile suggests deposition occurred along a well-oxygenated, low-relief ramp. 3D depositional facies models show Buda Formation evolution through time in Central Texas, and provide insights into reservoir evaluation and development.

Originally published as: Zhang, X., and M. C. Pope, 2017, High-resolution facies analysis and depositional architecture of the Buda Formation in Central Texas: Gulf Coast Association of Geological Societies Transactions, v. 67, p. 665.